



Ministério da Saúde
Instituto Nacional de Câncer
Coordenação de Ensino
Aperfeiçoamento nos Moldes Fellow em Radiologia Mamária

ANA LÉA BURNIER GANIMI COSTA

**Dermatofibrosarcoma Protuberans Simulating Primary
Breast Tumor**

RIO DE JANEIRO

2023

ANA LÉA BURNIER GANIMI COSTA

**Dermatofibrosarcoma Protuberans Simulating Primary
Breast Tumor**

Trabalho de Conclusão de Curso
apresentado ao Instituto Nacional
de Câncer como requisito parcial
para a conclusão do Curso de
Aperfeiçoamento nos moldes
Fellow em Radiologia Mamária.

Orientadora: Dra. Marcele França
Barreto Côrtes

Rio de Janeiro

2023

ANA LÉA BURNIER GANIMI COSTA

**Dermatofibrosarcoma Protuberans Simulating Primary
Breast Tumor**

Trabalho de Conclusão de Curso apresentado ao Instituto Nacional de Câncer
como requisito parcial para a conclusão do Curso de Aperfeiçoamento nos
Moldes Fellow em Radiologia Mamária

Aprovado em: 14 de outubro de 2023.

Banca examinadora:

Marcele França Barreto Côrtes

Dra. Marcelle França Barreto Côrtes

Paula Medina Gomes Maciel Curi Bonotto

Dra. Paula Medina Gomes Maciel Curi Bonotto

Érica Endo

Dra. Érica Endo

Rio de Janeiro

2023



DERMATOFIBROSARCOMA PROTUBERANS SIMULATING PRIMARY BREAST TUMOR

Ana Léa Burnier Ganimi Costa
Carolina Fernandes Ramalho
Giovana Ortiz D'Ávila
Marcele França Barreto Cortês
Thaís Salgado Monnerat
Camila de Carvalho Figueiredo
Érica Endo
Paula Medina Maciel Gomes Curi Bonotto

HOSPITAL NACIONAL DE CÂNCER - INCA 3

INTRODUCTION

- Dermatofibrosarcoma protuberans (DFSP) is a rare, slow-growing mesenchymal neoplasm of the dermis and subcutaneous tissues and tends to be locally aggressive.
- DFSP is characterized as low to intermediate grade neoplasm with high recurrence rate but rarely metastasises.

INTRODUCTION

- ▣ DFSP commonly involves the trunk and extremities, and very rarely the breast skin, mimicking a primary breast tumor with few reported cases in the literature.
- ▣ DFSP of the breast presents with a slow growing pattern, initially in the dermis and then invading the subcutaneous tissue.

DESCRIPTION

- ▣ We report an 18-year old female patient who was referred to our service, with a palpable mass growing in left breast.
- ▣ Assessment of the breasts with mammography revealed an irregular, indistinct, equal density mass in union of lower quadrants in left breast.
- ▣ On ultrasound, the lesion presents as oval, indistinct, complex cystic and solid mass, with vessels in rim. It was categorised as breast imaging reporting and data system (BIRADS) 4.

MAMMOGRAPHY

MLO

CC




Fig.1A

Fig.1B

Fig.1A and 1B: Mediolateral oblique and craniocaudal mammogram respectively, shows an irregular, indistinct, equal density mass in the union of lower quadrants in left breast, marked by metallic BB marker on skin (arrow).

ULTRASOUND




Fig.2A

Fig.2B

Fig.2C

Fig.2D

Fig. 2A and 2B: Ultrasound shows an oval, indistinct, complex cystic and solid mass (arrow).
Fig. 2C: Colour Doppler shows vessels in rim of the lesion (arrow).
Fig. 2D: Core biopsy of the lesion (arrow).

DESCRIPTION

- The patient underwent core biopsy of the lesion in left breast and the histopathological diagnosis was dermatofibrosarcoma protuberans.
- A local excision in a second time, confirmed the histopathological diagnosis.

DISCUSSION


- DFSP is a rare sarcoma of the dermis and subcutaneous tissue that has an estimated incidence of 0.8–5 cases per one million per year and accounts for less than 6% of all sarcomas.
- It commonly occurs on the trunk (50–60%), followed by the limbs (25%) and the head and neck area (10–15%). Involvement of the breast by DFSP is extremely rare, with only a few cases reported in the literature

DISCUSSION

- It usually affects patients aged between 20–50 years.
- DFSP has a high tendency of local recurrence and a low propensity for metastasis.
- Although rare, distant metastasis, especially in the lungs and lymph nodes, has been reported.

DISCUSSION


- Mammography: breast DFSP appears as a skin-based or intra-mammary located well-circumscribed oval-shaped lesion with no microcalcification nor speculation. These findings are not specific and hence, breast DFSP can mimic a benign breast lesion such as fibroadenoma, sebaceous cyst or an abscess. (Fig.3A e 3B)



Al-Farsi, Abeer et al., Dermatofibrosarcoma Protuberans Mimicking Primary Breast Neoplasm: A case report and literature review. (Accessed on 06 Jul 2023) DOI: 10.18295/squmj.2020.20.03.019

DISCUSSION

- Ultrasonographically: breast DFSP appears as a heterogeneously hyperechoic solid mass centered in subcutaneous tissue, with overlying hypoechoic region in the dermis with mild lobulation and increased vascularity on colour Doppler. Increased vascularity can be used to differentiate DFSP from benign breast lesions that usually appear as well-circumscribed lesions such as sebaceous cysts and liver abscesses. (Fig.4A e 4B)



Gilcrease-Garcia B, Dermatofibrosarcoma protuberans. Case study, Radiopaedia.org (Accessed on 06 Jul 2023) <https://doi.org/10.53347/rID-64660>

DISCUSSION

- Breast skin DFSP is a rare entity that can mimic a primary breast tumour.
- Awareness of this rare lesion and its typical imaging features will help to avoid the misdiagnosis of DFSP with a benign breast lesion and suggest a pre-operative diagnosis.

REFERENCES

- 1- Al-Farsi, A., Al-Brashdi, A., Al-Salhi, S., Al-Rahbi, N., Al-Rahbi, S., Al-Masqari, M., Al-Manji, Z., & Al-Umairi, R. S. (2020). Dermatofibrosarcoma Protuberans Mimicking Primary Breast Neoplasm: A case report and literature review. Sultan Qaboos University Medical Journal, 20(3), e368–371.
- 2- Kim H. Dermatofibrosarcoma protuberans in the breast: Case report. Int J Surg Case Rep. 2021 Apr;81:105773. doi: 10.1016/j.ijscr.2021.105773. Epub 2021 Mar 13. PMID: 33773373; PMCID: PMC8024653.
- 3- Kumar D, Vallacha A. Dermatofibrosarcoma Protuberans of Breast. J Coll Physicians Surg Pak. 2018 Aug;28(8):645-647. doi: 10.29271/jcpsp.2018.08.645. PMID: 30060798.
- 4- Dimas D, Boutas I, Potiris A, Koufopoulos N, Balalis D, Sitara K, Danglis F, Misitzis I. Dermatofibrosarcoma protuberans of the breast: A case study. Mol Clin Oncol. 2021 Mar;14(3):50. doi: 10.3892/mco.2021.2212. Epub 2021 Jan 20. PMID: 33604040; PMCID: PMC7849069.
- 5- Ferreira ALS, Souza JA, Gibbons IL, Lellis RF, Oliveira VS, de Sousa Rosas CH, Guatelli CS, Bitencourt AGV. Clinical and imaging features of dermatofibrosarcoma protuberans in the breast. Breast J. 2021 Jan;27(1):92-94. doi: 10.1111/tbj.14104. Epub 2020 Nov 27. PMID: 33246353.
- 6- Gilcrease-Garcia B, Dermatofibrosarcoma protuberans. Case study, Radiopaedia.org (Accessed on 06 Jul 2023) <https://doi.org/10.53347/rID-64660>.

THANK YOU



Email: analeaburnier@hotmail.com



Certificado

Declaramos que

*ANA LEA BURNIER GANIMI COSTA; CAROLINA FERNANDES RAMALHO;
GIOVANA ORTIZ D'ÁVILA; MARCELE FRANÇA BARRETO CÔRTEZ; THAIS
SALGADO MONNERAT; CAMILA DE CARVALHO FIGUEIREDO; ERICA
ENDO; PAULA MEDINA MACIEL GOMES CURI BONOTTO*

teve o trabalho intitulado "**DERMATOFIBROSARCOMA
PROTUBERANS SIMULATING PRIMARY BREAST TUMOR**"
aprovado e publicado em formato Pôster Eletrônico no 52º Congresso
Brasileiro de Radiologia e Diagnóstico por Imagem (CBR23).

São Paulo, 14 de outubro de 2023

Validação Online



Código: kuxY4gt4y3

Cibele Alves de Carvalho
Dra. Cibele Carvalho
Presidente do CBR

Ronaldo Hueb Baroni
Dr. Ronaldo Hueb Baroni
Diretor Científico do CBR